



Exploring Heterozygosity-Survival Correlations in a Wild Songbird Population: Contrasting Effects between Juvenile and Adult Stages

David Canal^{1*}, David Serrano², Jaime Potti¹

¹ Doñana Biological Station – CSIC, Department of Evolutionary Ecology, Sevilla, Spain, ² Doñana Biological Station – CSIC, Department of Conservation Biology, Sevilla, Spain

Abstract

The relationship between genetic diversity and fitness, a major issue in evolutionary and conservation biology, is expected to be stronger in traits affected by many loci and those directly influencing fitness. Here we explore the influence of heterozygosity measured at 15 neutral markers on individual survival, one of the most important parameters determining individual fitness. We followed individual survival up to recruitment and during subsequent adult life of 863 fledgling pied flycatchers born in two consecutive breeding seasons. Mark-recapture analyses showed that individual heterozygosity did not influence juvenile or adult survival. In contrast, the genetic relatedness of parents was negatively associated with the offspring's survival during the adult life, but this effect was not apparent in the juvenile (from fledgling to recruitment) stage. Stochastic factors experienced during the first year of life in this long-distance migratory species may have swamped a relationship between heterozygosity and survival up to recruitment.

Citation: Canal D, Serrano D, Potti J (2014) Exploring Heterozygosity-Survival Correlations in a Wild Songbird Population: Contrasting Effects between Juvenile and Adult Stages. PLoS ONE 9(8): e105020. doi:10.1371/journal.pone.0105020

Editor: Norman Johnson, University of Massachusetts, United States of America

Received: March 10, 2014; **Accepted:** June 3, 2014; **Published:** August 14, 2014

Copyright: © 2014 Canal et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are freely available. The data are publicly accessible at: <http://digital.csic.es/handle/10261/10044>.

Funding: This work was supported by projects PAC05-006-2, CGL2009-10652 and CGL2011-29694 from the Spanish Government. DC was supported by a grant from the Ministry of Education and Science (I3P-BDP2005), and also thanks Paola Laiolo for financial support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: davidcanal@ebd.csic.es

Introduction

Mating between related individuals may entail fitness costs to their descendents (e.g. [1–3]). The relationship between genetic diversity and fitness has therefore received much attention due to its potential importance in animal production, conservation and evolutionary biology (review in [4–9]). The study of this relationship in natural populations has traditionally been complex due to the difficulty of generating pedigrees to estimate individual coancestry [4]. With the expansion of molecular techniques in the last decades, however, an alternative approach based on the relationship between genetic diversity measured at a set of loci and traits related to fitness (heterozygosity-fitness correlation, HFC) has become widespread in the literature (review in [5,6,10]).

Positive HFCs, the association most commonly found in the literature [6], may arise mainly through three mechanisms [9,11]. Under the “direct hypothesis” heterozygous individuals have higher fitness due to overdominance of the typed loci. This mechanism may be important when heterozygosity is measured with allozymes or at the major histocompatibility complex (MHC) loci, but does not explain HFC in studies using microsatellites which, aside from exceptions (e.g. [12]), are considered neutral loci [13]. When employing microsatellites, two alternative hypotheses have been proposed to explain HFCs. First, the “local effect hypothesis” predicts that HFCs arise indirectly because the typed loci are linked to functional loci influencing fitness. Genetic drift,

migration and selection generate linkage favoring the detection of local effects [3,10]. Although linkage is expected to be low and rapidly eroded by recombination in natural populations [3,10], previous work has shown that relatively high levels of linkage may exist [14,15] and be maintained after more than 800 generations following a bottleneck in natural populations [14]. Second, under the “general effect hypothesis”, heterozygosity measured at multiple loci (MLH) would reflect heterozygosity across the genome. In such cases, homozygous individuals suffer fitness costs due to their higher likelihood of expression of deleterious recessive alleles (inbreeding depression; [1,2]). This mechanism is expected to arise under random mating, in populations with genetic drift, population admixture or suffering from recent bottlenecks, or in large populations where consanguineous matings occur. Populations under these conditions exhibit a large variance in inbreeding values, which determine the strength of the relationship between MLH and fitness [16,17].

Evidence of HFCs is widespread but also inconsistent in the literature (e.g. [18–23]). In general, HFCs are weak signals explaining no more than 3.6% of the variance in fitness [6]. Nevertheless, the magnitude of HFC may depend on the characteristics of the population (see above), the traits under scrutiny, and the environmental conditions that individuals experience [10,17,24]. HFCs have been commonly explored in behavioral (e.g. post-fledgling dispersal: [25]) or morphological

traits (e.g. body size: [26]; attractiveness: [27]), which are often under stabilizing selection [28]. However, evolutionary theory predicts [28], and empirical work confirms [5,6], greater HFCs in fitness-related traits under directional selection, as is the case of life history traits (e.g. fecundity, lifetime reproductive success, survival; [10]). This is because such traits are affected by multiple loci susceptible of deleterious recessive mutations [10]. Survival is one of the most important factors determining individual fitness and evidence of heterozygosity-survival correlations (HSC, hereafter) is common [19,29,30]. In birds, there is increasing information on HSC during early life (embryonic and nestling stages, or up to recruitment: e.g. [6,18,31]). Nevertheless, in the wild, HSC has only been explored beyond those stages, to our knowledge, in a Seychelles warbler (*Acrocephalus sechellensis*; [32,33]) and a blue tit (*Cyanistes caeruleus*; [12]) population.

HFCs are expected to decrease with age since differences in survival are greatest in early life [11]. Comprehensive studies across the lifespan of individuals, however, are crucial to the understanding of the mechanisms underlying HFCs since, for example, inbreeding effects may be underestimated [34,35] or even undetectable [36] when analyzed at a single stage. In addition, MLH effects may be negative early in life but positive during adult life [12]. In combination with age, the role of genetic diversity on fitness may also be sensitive to the environmental conditions that individuals experience [4,24]. Accordingly, recent studies highlight that, as a consequence of context-dependence, the magnitude of HFC may be inconsistent across years [37] or undetectable under favorable environmental conditions [33,38], i.e. if variance in the measured trait is affected by the environment, HFCs will be more easily detected in periods when environmental conditions cause large variation in the trait.

Here, we investigated the relationship between heterozygosity and individual survival across different life stages in a population of pied flycatchers (*Ficedula hypoleuca*), a long-distance passerine migrant. We followed individuals from two consecutive cohorts throughout their entire lifespan and explore individual survival using capture-recapture methods. This modeling framework is a more robust approach than generalized linear models when exploring effects on survival in open populations because it accounts for resighting probabilities [39,40]. Specifically, we explored whether: i) variation in survival was influenced by individual heterozygosity (MLH and/or single loci heterozygosity) or ii) by the genetic similarity or heterozygosity of an individual's parents, and iii) MLH and/or the magnitude of HFC changed across lifetime [11]. Thus, this study provides a suitable opportunity to investigate the influence of individual heterozygosity on survival in a wild population.

Materials and Methods

Ethics Statement

We hereby confirm that all data came from authorized monitoring of a natural population of pied flycatchers *Ficedula hypoleuca* studied in La Hiruela (province of Madrid, Spain, coordinates 41° 04' N, 3° 27' E) since 1984. Such long-term study requires birds being subjected to minimal disturbance and trying to keep animal suffering to a minimum. Manipulation was restricted to blood extraction from the brachial vein and all birds recovered well and were not released until we assessed their welfare. All fieldwork was conducted with the required permits for capture, ringing and blood sampling of birds by Consejería de Medio Ambiente, Comunidad de Madrid and Delegación de Medio Ambiente, Junta de Castilla-La Mancha.

Field work and general procedures

The study was carried out with individuals born in the breeding season of 2005 and 2006 as part of a long-term study of pied flycatchers in central Spain (e.g., [41,42]). The study area consists of two plots separated by 1.3 km, including 236 nest-boxes. Field protocols have been described in detail elsewhere [41]. Briefly, all nests were regularly checked every three days before the onset of egg laying and on a daily basis around hatching to ascertain laying date, clutch size, hatching date and number of fledglings. Parent birds were captured with a nest-box trap while feeding eight day-old nestlings. They were weighed, measured and individually marked with a numbered metal band and a unique combination of colored bands. Fledglings were banded, measured and weighed at 13 days of age. Blood samples were taken from all fledglings by puncturing the brachial vein and stored in absolute ethanol. Sex determination was carried out by PCR amplification of the CHD gene using the primers 2917 (forward) and 3088 (reverse; [43]). Molecular sexing was always fully consistent with the sex of recruited individuals.

Apparent survival of individual offspring was assessed through an extensive effort of marking, recapturing and resighting of color-banded birds in all subsequent breeding seasons until 2013. This population has high natal philopatry with a mean of 13% of recruitment of locally born birds, which is the highest recruitment rate found for the species [42,44,45].

Molecular methods

Our data set comprised 863 individuals born in 2005 ($n = 235$) and 2006 ($n = 628$), of which 101 recruited as breeders. Fledglings were genotyped at 15 polymorphic microsatellite loci: *f3-60*, *f1-25* [46], *fhy 216*, *fhy 237*, *fhy 301*, *fhy 304*, *fhy 310*, *fhy 329*, *fhy 339*, *fhy 356*, *fhy 361*, *fhy 401*, *fhy 444*, *fhy 466* and *fhy 236* [47]. Most individuals ($n = 835$; 96%) were genotyped at all loci, whereas genotyping failed for 23 individuals (2.6%) at one locus and for 5 individuals (0.6%) at two loci, respectively. Five individuals (0.6%) were discarded for further analyses since they were not genotyped for 4 or more loci.

Tests for Hardy-Weinberg equilibrium and linkage disequilibrium were done using the genotypes from the adult population of each study year with program Genepop 4.0 [48]. We performed a search in the zebra finch (*Taeniopygia guttata*) genome [49] to find the chromosome location of the loci used. A BLAT and BLAST search were run in UCSC (<http://genome.ucsc.edu/>) and ENSEMBL (http://www.ensembl.org/Taeniopygia_guttata/bblastview) browsers, respectively, to confirm the locations of the sequences. The best matched sequence was selected on the basis of both the lowest *E-value* and highest score. The "contig view" option in ENSEMBL was used to locate the nearest gene to the best matching sequence.

The set of markers used allowed us to identify cases of extra pair paternity and correct for it when exploring a paternal effect on fledgling survival (see below). Parentage analyses were carried out on CERVUS 3.0 [50] using a maximum likelihood method (see [41,51] for details). We considered a given male as the sire when he had a LOD (natural logarithm of the likelihood) score with a fledgling higher than the critical value requested for assignments at 95% confidence level (critical value is computed by CERVUS through parentage analyses simulations).

Estimation of heterozygosity and identity disequilibrium

Multilocus individual heterozygosity and allele frequencies were calculated with the Excel macro Cernicalin [52]. Cernicalin calculates three metrics: observed homozygosity per individual (HO), Internal relatedness (IR) and homozygosity by loci (HL).

The three metrics were highly correlated ($n = 863$, all $r > 0.97$, $p < 0.001$) and results did not vary among metrics (not shown). Analyses with HL are reported here because HL correlates better with genome-wide homozygosity and inbreeding in open populations than do other metrics [52]. HL varies between 0, when all loci in the individual are heterozygous, and 1, when all loci are homozygous. Homozygosity at a single locus (SHL) was coded as “0” for heterozygous status and “1” for homozygous. Pairwise relatedness coefficients of adults pairs (a proxy for inbreeding) were calculated through maximum likelihood estimation with software ML-RELATE [53].

Identity disequilibrium, the positive correlation between heterozygosity across loci, which is expected when HL is related to wide-genome heterozygosity, was calculated as g_2 in the program RMES [54]. Genotypes were resampled 1000 times to test if g_2 differed significantly from zero.

Capture-Recapture models

We used standard capture-mark-recapture (CMR) models implemented in software MARK 6.0 [55] to estimate apparent survival (ϕ) and recapture (p) probabilities [39] with maximum likelihood techniques (e.g. [56]). In particular, an extension of the Cormack-Jolly-Seber model including age and group effects was used [39]. By estimating (and correcting for) recapture probabilities, these models provide less-biased estimates of true survival probabilities than return rates, especially when site propensity and/or true detection are low (e.g. [57,58]). The recapture probability is the probability that a marked individual that is alive and present in the study area at sampling occasion t is captured (or observed) at sampling occasion t . The survival probability is the probability that a marked individual alive at sampling occasion t survived and has not permanently emigrated between sampling occasions t and $t+1$. These models make a number of assumptions to fit the available capture-recapture data. We tested the goodness of fit (GOF) of our data using U-CARE 2.3.2 [59].

In a first step, we tried to obtain a good fitting model in which potential genetic effects were not considered. Based on our previous knowledge of the species, four factors that could affect survival and recapture probabilities were included: time, sex, age (fledgling, 1-yr, 2-yr and older) and cohort (2005 versus 2006). We began with a global model in which all parameters were different (denoted ϕ_{s*t*c} , p_{s*t*c} , where s = sex, t = time and c = cohort). Given that age and time are identical within a given cohort, the three factors cannot be tested simultaneously, so we built plausible pairwise combinations of the three variables. We tried to constrain these initial models by building a set of candidate models with fewer parameters that had a reasonable biological justification (i.e. reducing the number of age-classes and assuming no differences among years and/or between sexes and cohorts). Once the most parsimonious models were identified, we added genetic data (genetic relatedness of parents and multilocus heterozygosity of fledglings and parents) as individual covariates to test their importance on apparent survival. Model selection was based on the Akaike's information criterion adjusted for small sample sizes (AICc) and Akaike weights (w_i , [60]). In addition, we used the analysis of deviance test (ANODEV, [61]) to compare the amount of deviance explained by a covariate.

To test for single locus effects on survival, we additionally built in a generalized linear modeling framework (GLMM with a binomial error distribution) i) a model including all single locus heterozygosities (SHL), and ii) a model including the individual HL, and compared them with a F-ratio test as suggested by Szulkin et al. [62]. In the model containing heterozygosities at all single loci, missing data at one locus were replaced by the sample

average at that locus [62]. The procedure was performed for both juvenile and adult survival regardless of whether or not general effects were found. In fact, significant SLH effects may exist even in absence of HL correlation as loci may have similar effects in opposite directions [22]. Sample sizes varied slightly across statistical analyses because all information was not always available for all individuals.

Results

General genetic parameters

Individual HL ranged from 0 to 0.56, with a mean (\pm SD) of $0.208 (\pm 0.102)$. HL did not differ between years (2005, $n = 234$: 0.198 ± 0.102 ; 2006, $n = 620$: 0.212 ± 0.102 , GLMM: $F_{1,666} = 1.10$, $p = 0.29$), sex (males, $n = 477$: 0.205 ± 0.103 ; females, $n = 377$: 0.212 ± 0.100 , GLMM: $F_{1,156} = 0.68$, $p = 0.40$) or among age groups (fledglings: 0.21 ± 0.1 ; yearlings: 0.20 ± 0.09 ; older individuals: 0.20 ± 0.09 ; $F_{1,156} = 0.22$, $p = 0.64$; interaction age*sex: $F_{1,154} = 0.05$, $p = 0.78$). Nestlings' HL and their parent relatedness were weakly but significantly correlated ($r = 0.32$; $p < 0.001$).

The search within the zebra finch putative genome showed that the markers we used were widespread throughout the passerine genome (Table 1). All loci conformed to Hardy-Weinberg equilibrium and no pair of loci showed significant linkage disequilibrium after Bonferroni correction. Identity disequilibrium was low and did not differ significantly from zero ($g_2 = 0.0006$, $sd = 0.0008$, $p = 0.17$).

Apparent survival

The U-CARE goodness of fit global test was highly significant ($\chi^2 = 36.93$, d.f. = 12, $p = 0.0002$). The specific test for transience was also significant ($Z = 2.99$, $p = 0.001$), while there was no statistical evidence for trap-dependence ($Z = -1.23$, $p = 0.22$). After suppressing the first occasion from the capture-recapture history, the GOF tests were no longer significant (Global test: $\chi^2 = 6.21$, d.f. = 8, $p = 0.62$; directional test for transience: $Z = 1.16$, $p = 0.12$), indicating that a true age effect was responsible for the lack of fit (i.e. the transient effect was associated with differential capture-recapture probability between juveniles and adults). We accounted for this in our models by estimating age-specific survival and recapture parameters.

The initial modelling process without genetic covariates revealed that two models received most support. The first model (ϕ_{a2} , p_{a3*s*c} , AICc = 1092.699, $K = 11$, $w_i = 0.398$, Deviance = 1070.40) showed that survival was different for juveniles and adults, while recapture varied with time, and between sexes and cohorts. The second top-ranked model (ϕ_{a2*s} , p_{a3*s*c} , AICc = 1094.559, Δ AICc = 1.859, $K = 13$, $w_i = 0.157$, Deviance = 1068.15) had the same structure but included a sex effect for apparent survival. The remaining models were separated by > 4 AICc points and thus considered poorly supported (Results not shown).

Table 2 shows the overall best models once the effect of genetic covariates was incorporated. The top-ranked model (model 1) indicated that relatedness of the parents affected offspring survival during adult life but not between fledgling and recruitment; parents with a higher relatedness coefficient raised fledglings that had lower survival probabilities throughout adulthood (Fig. 1). The covariate in this model has a highly significant effect (ANODEV test: $F_{1,17} = 62.86$, $p < 0.0001$) and explained 78.7% of the survival variability. The second top-ranked model included also an effect on juvenile survival, but the examination of 95% confidence intervals for the estimate of the effect indicated that this parameter is confounding [60,63]. Indeed, the model including

Table 1. Characteristics of the microsatellite loci used in the study.

Locus	A	Ho	He	Chromosome: Start (Bp)	ID nearest gene*	Distance (Bp)
f1-25	7	0.738	0.7554	20/10.407.082	07700	exon ⁺
f3-60	35	0.9543	0.961	9/14.479.023	09067	27.807
Fhy 216	8	0.518	0.521	1a/63.001.644	11914	11.2742
Fhy 236	25	0.896	0.87	20/13.791.266	08550	11.5271
Fhy 237	6	0.399	0.392	3/7.930.679	02614	15.700
Fhy 301	14	0.856	0.884	2/92.250.591	07400	150.859
Fhy 304	10	0.79	0.803	4_random/2.365.290	15203	763.363
Fhy 310	13	0.872	0.864	2/92.250.591	15081	2.257
Fhy 329	8	0.682	0.672	3/49.130.923	10789	68.805
Fhy 339	12	0.831	0.83	1/95.843.488	13407	3.561
Fhy 356	12	0.833	0.856	1a/6.627.591	02324	29.231
Fhy 361	7	0.549	0.518	2/29.361.136	01714	195.077
Fhy 401	13	0.795	0.788	Un/52.369.084	06481	696.145
Fhy 444	14	0.8757	0.8816	1/12.170.793	07097	293.505
Fhy 466	12	0.8362	0.8438	7/21.099.737	10012	5.274

Number of alleles (A), observed (Ho) and expected (He) heterozygosities, chromosome location, and distance to the nearest gene (in base pairs; Bp) according to their position in the zebra finch genome are shown.

*Last digits of the gene's ID in ENSEMBL. Prefix: ENSTGUG000000.

⁺located in a exon of a solute carrier organic anion transporter gene.

doi:10.1371/journal.pone.0105020.t001

survival from fledgling to recruitment only (model 13) received little support. In any case, models incorporating the effect of relatedness of the parents on survival had a cumulative weight of 0.977, being 43 times better supported than models without this effect. Models including the father's (models 7, 9 and 11) or the mother's (models 10 and 15) multilocus heterozygosity were hardly

supported, as was a model with multilocus fledgling heterozygosity (model 14) or models without any covariate (models 8 and 12).

The variance explained did not differ between a model including heterozygosity at all single loci and a model including HL for juvenile ($F_{27,506} = 1.20$, $p = 0.22$) and adult ($F_{27,506} = 0.94$, $p = 0.54$) ages suggesting an absence of single loci effects on survival [62].

Table 2. Top supported models (90% confidence set) for apparent survival (ϕ) and recapture (p) of pied flycatchers.

No.	Model	AICc	$\Delta AICc$	K	w_i	Deviance
1	$\phi_{a2(2+ : REL), p_{a3*s*c}}$	1082.7384	0	12	0.46579	1058.3874
2	$\phi_{a2*REL, p_{a3*s*c}}$	1083.0954	0.357	13	0.38964	1056.6855
3	$\phi_{a2(2+ : REL), p_{a3*s}}$	1086.9571	4.2187	9	0.05651	1068.7553
4	$\phi_{a2*REL, p_{a3*s}}$	1087.476	4.7376	10	0.04359	1067.2291
5	$\phi_{(a2*c*REL) + sr, p_{a3*s}}$	1089.8896	7.1512	14	0.01304	1061.4161
6	$\phi_{a2+ REL, p_{a3*s*c}}$	1091.0027	8.2643	12	0.00748	1066.6517
7	$\phi_{a2(1 \text{ yr: HL}_F), p_{a3*s*c}}$	1092.3524	9.614	12	0.00381	1068.0014
8	$\phi_{a2, p_{a3*s*c}}$	1092.6996	9.9612	11	0.0032	1070.403
9	$\phi_{a2*HL_F, p_{a3*s*c}}$	1092.7067	9.9683	13	0.00319	1066.2968
10	$\phi_{a2(1 \text{ yr: HL}_M), p_{a3*s*c}}$	1093.4256	10.6872	12	0.00223	1069.0746
11	$\phi_{(a2*HL_F) + sr, p_{a3*s*c}}$	1094.3387	11.6003	14	0.00141	1065.8652
12	$\phi_{a2*sr, p_{a3*s*c}}$	1094.5591	11.8207	13	0.00126	1068.1492
13	$\phi_{a2(1 \text{ yr: REL}), p_{a3*s*c}}$	1094.6914	11.953	12	0.00118	1070.3404
14	$\phi_{a2(1 \text{ yr: HL}), p_{a3*s*c}}$	1094.7269	11.9885	12	0.00116	1070.3759
15	$\phi_{(a2*HL_M) + sr, p_{a3*s*c}}$	1095.0261	12.2877	14	0.001	1066.5526

Shown are Akaike information criteria corrected for small sample sizes (AICc), difference in AICc with the top-ranked model ($\Delta AICc$), number of estimable parameters (K), normalized Akaike weights (w_i), and deviance. Subscripts denote age (a2 and a3 corresponding to two and three age-classes respectively), sex (s) and cohort (c). Covariates for heterozygosity: fledgling (HL), father (HL_F), mother (HL_M) and genetic relatedness of the parents (REL). '1 yr' and '2+:' denote an effect only present from fledgling to first-year and from first-year onwards, respectively. Symbol '*' denotes interaction, and symbol '+' additive effects.

doi:10.1371/journal.pone.0105020.t002




Figure 1. Apparent survival probability of adults (from their first-year onwards) in relation to the pairwise genetic relatedness of their parents. Low values indicate lower relatedness (e.g. relatedness between parent-offspring or full siblings is 0.5). Dashed lines denote 95% confidence limits around the predicted linear trend.
doi:10.1371/journal.pone.0105020.g001

Discussion

Using a wild population of a migratory songbird we explored the relationship between HL and apparent survival at different life stages. Contrary to our expectations, HL was not related to survival either in the juvenile or in subsequent adult life stages. Remarkably, however, the best ranked model in the CMR analyses indicated that the genetic relatedness of the parents (a proxy for inbreeding) negatively influenced offspring survival during adult life, but such effect was not apparent from fledgling to recruitment. Although true survival and permanent emigration are confounded in Cormack-Jolly-Seber models [58], our findings showing an effect during the adult stage suggest an actual relationship between the genetic relatedness of the parents and mortality. First, adults show much higher site fidelity rates and much shorter dispersal distances than juveniles, and our study area was large enough to encompass most between-years movements of breeding individuals (mean breeding dispersal distance = 150 m; [64–66]). Second, mean apparent survival probabilities of adult birds (55%, see below), were higher than the highest value reported for other populations of the species (e.g. [67–69]).

As expected from theory, positive relationships between heterozygosity and survival, an important component of fitness, have been commonly reported in a variety of taxa (amphibians: [38]; mammals: [19,29]; birds: [18,31]). Most of these studies have explored survival in early life stages (i.e. embryo, juvenile or up to recruitment) likely due to the logistical constraints in obtaining longitudinal data across the lifespan of individuals in open populations, and also because the magnitude of HFC is expected to decrease with age [11,20,70]. However, measuring HFC at a single life-stage and/or monitoring individuals along only a fraction of their lifespans could cause bias in the reported results. In fact, evidence of HFCs with increasing age has been also reported, with old individuals being more sensitive to environmental variation than young ones [71]. In addition, even opposite effects of heterozygosity have been detected at different life stages with negative and positive effects in early and late life, respectively [12,36,72,73]. Similarly, an effect of the parent's relatedness on the offspring survival only became apparent in our population after recruitment whereas it was undetectable from fledgling to recruitment (see below). Pied flycatchers are long-distance migrants and, given that HFCs explain on average 1% of fitness variation [6] and fledgling recruitment is low (on average, 22%

compared to 55% of adult survival in the study years, as estimated from model 8), stochastic factors operating during migration and/or soon after fledging (e.g. severe adverse conditions and/or high rates of predation) may have overridden any detectable effect of heterozygosity on individual survival up to recruitment. The contrasting effects at different life stages found here add to those in the literature and highlight the need of additional work concerning HSC beyond juvenile phases to clarify its impact in wild populations.

Heterozygosity of the genetic parents (as a specific parental effect; see e.g. [74] or their genetic relatedness have been reported to affect offspring survival. In blue tits, parental HL was positively related to recruitment rate [12] whereas in Seychelles warblers, fledgling survival was influenced by the heterozygosity of both the genetic father [32]; but see [33]) and mother [33]. Our capture-recapture analyses suggest that pairs formed by genetically similar birds raised individuals with low survival prospects after recruiting as breeders. Further, models including an effect of the father heterozygosity were in general better supported than models without any covariate leading support to the idea that there is a genetic effect of the parents on offspring survival (although the effect of mother heterozygosity was less informative). In this regard, the weak association between the offspring heterozygosity and the relatedness of their parents found here could partly explain the lack of a direct relationship between individual HL and survival in our population. Due to the nature of the markers employed (highly polymorphic microsatellites), the probability of having a high HL (i.e. homozygous individuals) may be low even with genetically similar parents, which may obscure a direct HSC.

None of the loci used appear to be linked to functional genes possibly influencing survival. Further, the lack of identity disequilibrium (estimated as g_2 ; see methods) suggests that the typed loci are unrelated to inbreeding at the genome level. However, HFCs can occur (and generally do occur; [62]) in the absence of identity disequilibrium: given that traits are usually influenced by many more loci than those typed, slight inbreeding effects are more easily detected on the phenotype (through HFC) than through correlations in the loci [62]. Thus, although a larger panel of markers should apparently report more precise estimates of inbreeding [17,75], this argument should not be used to invalidate HFC works [62]. In fact, a recent study of a zebra finch population (with low inbreeding variance) has challenged this view by showing that a panel of 11 microsatellites (the mean microsatellite number used in HFC studies; [6]) located across the genome is as informative as a panel of 1359 SNP markers or a

5th generation pedigree [23]. In the same line, Taylor et al. [76] reported a significant correlation between a pedigree-based inbreeding (f) and a panel of 13 microsatellites suggesting that a small set of markers is not necessarily uninformative on individual's f even in wild, outbred populations. In open populations, immigration may generate linkage disequilibrium which, together with random mating, cause identity disequilibrium (i.e. inbreeding; [10]). Unfortunately, we could not measure the degree of the relationship between HL and a pedigree-based inbreeding in this study. The rate of immigration to the population is high and given that reliable inbreeding values must be based on at least two generations (i.e. grandparents-grandsons), we knew the identity of nestlings and grandparents in only 15 broods, with the inbreeding value being zero in all cases. An unsurprising result, since mating between first-order relatives seems to be infrequent in natural populations (around 3%; [77]).

More than a decade ago, Keller and Walker [4] highlighted the need of studies exploring the interaction between genetics, environment and fitness. Today, this type of study is still very scarce. Recent work has highlighted the role of heterozygosity on fitness even in large natural populations with apparent absence of inbreeding [21], which could be determined by environmental conditions [33,37] or, as in the present work, detected at late stages of life. Hence, we emphasize that studies exploring HFCs at different life stages, in populations with different demographic histories and under variable environmental conditions are required to increase our knowledge on the causes of HFCs.

Supporting Information

Database S1

(XLSX)

Acknowledgments

We would like to thank Inés Valencia and Carlos Camacho for their dedicated assistance in the field. We are grateful to the associate editor Dr. Norman Johnson and one anonymous reviewer for constructive comments improving this work.

Author Contributions

Conceived and designed the experiments: DC JP. Performed the experiments: DC JP. Analyzed the data: DC JP DS. Contributed reagents/materials/analysis tools: DC JP DS. Wrote the paper: DC JP DS.

References

- Charlesworth B, Charlesworth D (1987) Inbreeding Depression And Its Evolutionary Consequences. *Annu Rev Ecol Syst* 18: 237–268.
- Charlesworth B, Charlesworth D (1999) The genetic basis of inbreeding depression. *Genet Res* 74: 329–340.
- Lynch M, Walsh B (1998) *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, editor Sunderland, Massachusetts.
- Keller L, Waller D (2002) Inbreeding effects in wild populations. *Trends Ecol Evol* 17: 19–23.
- Coltman D, Slate J (2003) Microsatellite measures of inbreeding: A meta-analysis. *Evolution* 57: 971–983.
- Chapman JR, Nakagawa S, Coltman DW, Slate J, Sheldon BC (2009) A quantitative review of heterozygosity-fitness correlations in animal populations. *Mol Ecol* 18: 2746–2765.
- Kempnaers B (2007) Mate choice and genetic quality: a review of the heterozygosity theory. *Adv study Behav* 37: 189–278.
- Hedrick P, Kalinowski S (2000) Inbreeding depression in conservation biology. *Annu Rev Ecol Syst* 31: 139–162.
- Hansson B, Westberg L (2002) On the correlation between heterozygosity and fitness in natural populations. *Mol Ecol* 11: 2467–2474.
- Szulkin M, Bierne N, David P (2010) Heterozygosity-fitness correlations: a time for reappraisal. *Evolution* 64: 1202–1217.
- David P (1998) Heterozygosity-fitness correlations: new perspectives on old problems. *Heredity* 80: 531–537.
- Olano-Marin J, Mueller JC, Kempnaers B (2011) Heterozygosity and survival in blue tits (*Cyanistes caeruleus*): contrasting effects of presumably functional and neutral loci. *Mol Ecol* 20: 4028–4041.
- Jarne P, Lagoda PJ (1996) Microsatellites, from molecules to populations and back. *Trends Ecol Evol* 11: 424–429.
- Reich DE, Cargill M, Bolk S, Ireland J, Sabeti PC, et al. (2001) Linkage disequilibrium in the human genome. *Nature* 411: 199–204.
- Dawson E, Abecasis R, Bumpstead S, Chen Y, Hunt S, et al. (2002) A first-generation linkage disequilibrium map of human chromosome 22. *Nature* 418: 544–548.
- Balloux F, Amos W, Coulson T (2004) Does heterozygosity estimate inbreeding in real populations? *Mol Ecol* 13: 3021–3031. A.
- Slate J, David P, Dodds KG, Veenivliet BA, Glass BC, et al. (2004) Understanding the relationship between the inbreeding coefficient and multi-locus heterozygosity: theoretical expectations and empirical data. *Heredity* 93: 255–265.
- Hansson B, Bensch S, Hasselquist D, Akesson M (2001) Microsatellite diversity predicts recruitment of sibling great reed warblers. *Proc R Soc B Biol Sci* 268: 1287–1291.

19. Acevedo-Whitehouse K, Spraker TR, Lyons E, Melin SR, Gulland F, et al. (2006) Contrasting effects of heterozygosity on survival and hookworm resistance in California sea lion pups. *Mol Ecol* 15: 1973–1982.
20. Rijks JM, Hoffman JL, Kuiken T, Osterhaus AD, Amos W (2008) Heterozygosity and lungworm burden in harbour seals (*Phoca vitulina*). *Heredity* 100: 587–593.
21. Wetzel DP, Stewart IRK, Westneat DF (2012) Heterozygosity predicts clutch and egg size but not plasticity in a house sparrow population with no evidence of inbreeding. *Mol Ecol* 21: 406–420.
22. Chapman JR, Sheldon BC (2011) Heterozygosity is unrelated to adult fitness measures in a large, noninbred population of great tits (*Parus major*). *J Evol Biol* 24: 1715–1726.
23. Forstmeier W, Schielzeth H, Mueller JC, Ellegren H, Kempenaers B (2012) Heterozygosity-fitness correlations in zebra finches: microsatellite markers can be better than their reputation. *Mol Ecol* 21: 3237–3249.
24. Armbruster P, Reed DH (2005) Inbreeding depression in benign and stressful environments. *Heredity* 95: 235–242.
25. Gillingham MAF, Cézilly F, Wattier R, Béchet A (2013) Evidence for an association between post-fledging dispersal and microsatellite multilocus heterozygosity in a large population of greater flamingos. *PLOS One* 8: e81118.
26. Ryder TB, Tori WP, Blake JG, Loiselle BA, Parker PG (2009) Mate choice for genetic quality: a test of the heterozygosity and compatibility hypotheses in a lek-breeding bird. *Behav Ecol* 21: 203–210.
27. Foerster K, Delhey K, Johnson A, Liffield JT, Kempenaers B (2003) Females increase offspring heterozygosity and fitness through extra-pair matings. *Nature* 425: 714–717.
28. Houle D, Morikawa B, Lynch M (1996) Comparing mutational variabilities. *Genetics* 143: 1467–1483.
29. Coulson TN, Pemberton JM, Albon SD, Beaumont M, Marshall TC, et al. (1998) Microsatellites reveal heterosis in red deer. *Proc R Soc B Biol Sci* 265: 489–495.
30. Da Silva A, Gaillard J-M, Yoccoz NG, Hewison AJM, Galan M, et al. (2009) Heterozygosity-fitness correlations revealed by neutral and candidate gene markers in roe deer from a long-term study. *Evolution* 63: 403–417.
31. Jensen H, Bremset EM, Ringsby TH, Saether B-E (2007) Multilocus heterozygosity and inbreeding depression in an insular house sparrow metapopulation. *Mol Ecol* 16: 4066–4078.
32. Richardson DS, Komdeur J, Burke T (2004) Inbreeding in the Seychelles warbler: environment-dependent maternal effects. *Evolution* 58: 2037–2048.
33. Brouwer L, Komdeur J, Richardson DS (2007) Heterozygosity-fitness correlations in a bottlenecked island species: a case study on the Seychelles warbler. *Mol Ecol* 16: 3134–3144.
34. Szulkin M, Garant D, McCleery RH, Sheldon BC (2007) Inbreeding depression along a life-history continuum in the great tit. *J Evol Biol* 20: 1531–1543.
35. Grueber CE, Laws RJ, Nakagawa S, Jamieson IG (2010) Inbreeding depression accumulation across life-history stages of the endangered Takahe. *Conserv Biol* 24: 1617–1625.
36. Von Hardenberg A, Bassano B, Festa-Bianchet M, Luikart G, Lanfranchi P, et al. (2007) Age-dependent genetic effects on a secondary sexual trait in male Alpine ibex, *Capra ibex*. *Mol Ecol* 16: 1969–1980.
37. Harrison XA, Bearhop S, Inger R, Colhoun K, Gudmundsson GA, et al. (2011) Heterozygosity-fitness correlations in a migratory bird: an analysis of inbreeding and single-locus effects. *Mol Ecol* 20: 4786–4795.
38. Lesbarrères D, Primmer CR, Laurila A, Merilä J (2005) Environmental and population dependency of genetic variability-fitness correlations in *Rana temporaria*. *Mol Ecol* 14: 311–323.
39. Lebreton J-D, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and testing biological hypotheses using marked animals: A unified approach with case studies. *Ecol Monogr* 62: 67–118.
40. Kalbfleisch JD, Prentice RL (2002) The statistical analysis of failure time data. 2nd edn. New York: John Wiley & Sons, Inc.
41. Canal D, Potti J, Dávila JA (2011) Male phenotype predicts extra-pair paternity in pied flycatchers. *Behaviour* 148: 691–712.
42. Potti J, Canal D, Serrano D (2013) Lifetime fitness and age-related female ornament signalling: evidence for survival and fecundity selection in the pied flycatcher. *J Evol Biol* 26: 1445–1457.
43. Ellegren H (1996) First gene on the avian W chromosome (CHD) provides a tag for universal sexing of non-ratite birds. *Proc R Soc B Biol Sci* 263: 1635–1641.
44. Potti J, Montalvo S (1991) Male arrival and female mate choice in Pied Flycatchers *Ficedula hypoleuca* in Central Spain. *Ornis Scand* 22: 45–54.
45. Lundberg A, Alatalo R (1992) The pied flycatcher. London: Poyser.
46. Molecular Ecology Resources Primer Development Consortium, Abercrombie LG, Anderson CM, Baldwin BG, Bang IC, et al. (2009) Permanent Genetic Resources added to Molecular Ecology Resources database 1 January 2009–30 April 2009. *Mol Ecol Resour* 9: 1375–1379.
47. Leder EH, Karaïskou N, Primmer CR (2008) Seventy new microsatellites for the pied flycatcher, *Ficedula hypoleuca* and amplification in other passerine birds. *Mol Ecol Resour* 8: 874–880.
48. Raymond M, Rousset F (1995) Genepop (version 1.2), population genetics software for exact tests and ecumenicism. *J Hered* 86: 248–249.
49. Balakrishnan CN, Ekblom R, Völker M, Westerdahl H, Godínez R, et al. (2010) Gene duplication and fragmentation in the zebra finch major histocompatibility complex. *BMC Biol* 8: 29.
50. Marshall T, Slate J, Bruuk L, Pemberton JM (1998) Statistical confidence for likelihood-based paternity. *Mol Ecol* 7: 639–655.
51. Canal D, Jovani R, Potti J (2012) Multiple mating opportunities boost protandry in a pied flycatcher population. *Behav Ecol Sociobiol* 66: 67–76.
52. Aparicio JM, Ortego J, Cordero PJ (2006) What should we weigh to estimate heterozygosity, alleles or loci? *Mol Ecol* 15: 4659–4665.
53. Kalinowski ST, Wagner AP, Taper ML (2006) ML-RELATE: a computer program for maximum likelihood estimation of relatedness and relationship. *Mol Ecol Notes* 6: 576–579.
54. David P, Pujol B, Viard F, Castella V, Goudet J (2007) Reliable selfing rate estimates from imperfect population genetic data. *Mol Ecol* 16: 2474–2487.
55. White GC, Burnham KP (1999) Program MARK: survival estimation from populations of marked animals. *Bird Study* 46: S120–S139.
56. Williams BK, Nichols JD, Conroy MJ (2002) Analysis and Management of Animal Populations. Academic, San Diego, California, USA.
57. Martin TE, Clobert J, Anderson DR (1995) Return rates in studies of life history evolution: Are biases large? *J Appl Stat* 22: 863–876.
58. Sandercock BK (2006) Estimation of Demographic Parameters from Live-Encounter Data: a Summary Review. *J Wildl Manage* 70: 1504–1520.
59. Choquet R, Lebreton J-D, Gimenez O, Reboulet A-M, Pradel R (2009) U-CARE: Utilities for performing goodness of fit tests and manipulating CAPTURE-REcapture data. *Ecography* 32: 1071–1074.
60. Burnham KP, Anderson DR (2002) Model selection and multimodel inference. A practical information-theoretic approach. Springer.
61. Skalski J, Hoffmann A, Smith S (1993) Marked Individuals in the Study of Bird Population. In: Testing the significance of individual- and cohort-level covariates in animal survival studies. J.D Lebreton & P.M North, editor. Birkhäuser-Verlag, Basle, Switzerland. pp.9–28.
62. Szulkin M, Bierne N, David P (2010) Heterozygosity-fitness correlations: a time for reappraisal. *Evolution* 64: 1202–1217.
63. Arnold TW (2010) Uninformative Parameters and Model Selection Using Akaike's Information Criterion. *J Wildl Manage* 74: 1175–1178.
64. Potti J, Montalvo S (1991) Return rate, age at first breeding and natal dispersal of pied flycatchers *Ficedula hypoleuca* in Central Spain. *Ardea* 79: 419–428.
65. Montalvo S, Potti J (1992) Breeding Dispersal in Spanish Pied Flycatchers *Ficedula hypoleuca*. *Ornis Scand* 23: 491.
66. Camacho C, Canal D, Potti J (2013) Nonrandom dispersal drives phenotypic divergence within a bird population. *Ecol Evol* 3: 4841–4848.
67. Huk T, Winkel W (2006) Polygyny and its fitness consequences for primary and secondary female pied flycatchers. *Proc Biol Sci* 273: 1681–1688.
68. Eeva T, Hakkarainen H, Laaksonen T, Lehikoinen E (2006) Environmental pollution has sex-dependent effects on local survival. *Biol Lett* 2: 298–300.
69. Chernetsov N, Sokolov LV, Kosarev V (2009) Local survival rates of Pied Flycatchers *Ficedula hypoleuca* depend on their immigration status. *Avian Ecol Behav*: 11–20.
70. Cohas A, Bonenfant C, Kempenaers B, Allainé D (2009) Age-specific effect of heterozygosity on survival in alpine marmots, *Marmota marmota*. *Mol Ecol* 18: 1491–1503.
71. Charlesworth B, Hughes KA (1996) Age-specific inbreeding depression and components of genetic variance in relation to the evolution of senescence. *Proc Natl Acad Sci U S A* 93: 6140–6145.
72. Olano-Marin J, Mueller JC, Kempenaers B (2011) Correlations between heterozygosity and reproductive success in the blue tit (*Cyanistes caeruleus*): an analysis of inbreeding and single locus effects. *Evolution* 65: 3175–3194.
73. Escobar JS, Nicot A, David P (2008) The different sources of variation in inbreeding depression, heterosis and outbreeding depression in a metapopulation of *Physa acuta*. *Genetics* 180: 1593–1608.
74. Price T (1998) Maternal and paternal effects in birds. In: Mousseau, T.A. and Fox CW, editor. Maternal Effects as Adaptations. Oxford University Press. pp.202–226.
75. Miller JM, Malenfant RM, David P, Davis CS, Poissant J, et al. (2013) Estimating genome-wide heterozygosity: effects of demographic history and marker type. *Heredity* 112: 240–247.
76. Taylor SS, Sardell RJ, Reid JM, Bucher T, Taylor NG, et al. (2010) Inbreeding coefficient and heterozygosity-fitness correlations in unhatched and hatched song sparrow nestmates. *Mol Ecol* 19: 4454–4461.
77. Ralls K, Harvey P, Lyles A (1986) Inbreeding in natural populations of birds and mammals. In: M.E. Soulé, editor. Conservation Biology. Sinauer Associates, Sunderland, MA. pp.35–56.